

### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

Kilometers

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

•A: Limited potential for groundwater flooding to occur
•B: Potential for groundwater flooding of property situated below ground level
•C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL LEVEL 1 SFRA


**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding

Index Number: ND\_01

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of Jeremy Benn Associates Ltd.







### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

Kilometers

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

•A: Limited potential for groundwater flooding to occur
•B: Potential for groundwater flooding of property situated below ground level
•C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL **LEVEL 1 SFRA**

**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding


Sheet No: 2 of 32


Index Number: ND\_02

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of Jeremy Benn Associates Ltd.









### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings.

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

- •A: Limited potential for groundwater flooding to occur
  •B: Potential for groundwater flooding of property situated below ground level
  •C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL LEVEL 1 SFRA

**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding


Sheet No: 4 of 32

Index Number: ND\_04

This document is the property of Jeremy Benn Associates Ltd. It shall not be losed to a third party, without the permission of eproduced in whole or in part, nor disc Jeremy Benn Associates Ltd.







North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at

Kilometers

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Jnauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.


### NORTH DORSET DISTRICT COUNCIL **LEVEL 1 SFRA**


**BGS** Susceptibility to Groundwater Flooding

Index Number: ND\_05

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of







### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

Kilometers

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Jnauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

- •A: Limited potential for groundwater flooding to occur
  •B: Potential for groundwater flooding of property situated below ground level
  •C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

### NORTH DORSET DISTRICT COUNCIL **LEVEL 1 SFRA**

**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding


Sheet No: 6 of 32

Index Number: ND\_06

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of Jeremy Benn Associates Ltd.







Limited potential for groundwater flooding to occur


Potential for groundwater flooding to occur at


Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Inauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

**BGS** Susceptibility to Groundwater Flooding

eproduced in whole or in part, nor disclosed to a third party, without the permission of





### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

ntains Environment Agency information © Environment Agency and/or database righ Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings.

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

- •A: Limited potential for groundwater flooding to occur
  •B: Potential for groundwater flooding of property situated below ground level
  •C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL **LEVEL 1 SFRA**

**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding


Sheet No: 8 of 32


Index Number: ND\_08


This document is the property of Jeremy Benn Associates Ltd. It shall not be osed to a third party, without the permission of Jeremy Benn Associates Ltd.











### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at surface

Kilometers

Environment Agency information © Environment Agency and/or database righ Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Inauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

- •A: Limited potential for groundwater flooding to occur •B: Potential for groundwater flooding of property situated below ground level •C: Potential for groundwater flooding to occur at surface

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL LEVEL 1 SFRA


**APPENDIX A** 


**BGS** Susceptibility to Groundwater Flooding

Index Number: ND\_11

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of Jeremy Benn Associates Ltd.







North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property

Potential for groundwater flooding to occur at

2 Kilometers

Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) nauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.


The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:


Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL

**BGS** Susceptibility to Groundwater Flooding

eproduced in whole or in part, nor disclosed to a third party, without the permission of





North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at

Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings.


The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.


The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:


Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.


# NORTH DORSET DISTRICT COUNCIL **LEVEL 1 SFRA**

This document is the property of Jeremy Benn Associates Ltd. It shall not be osed to a third party, without the permission of









### Legend

North Dorset District Council Boundary

Limited potential for groundwater flooding to occur

Potential for groundwater flooding of property situated below ground level

Potential for groundwater flooding to occur at

Contains Environment Agency information © Environment Agency and/or database right Reproduced from Ordnance Survey mapping with the permission of Ordnance Survey on behalf of the Controller of Her Majesty's Stationery Office. © Crown copyright. (2017) Inauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings

### **Technical Note**

The BGS Susceptibility to Groundwater Flooding dataset is a national dataset covering England, Wales and Scotland. Based on geological and hydrogeological information, the digital data can be used to identify areas where geological conditions could enable groundwater flooding to occur and where groundwater may come close to the ground surface. It is based on a 50m square grid.

The dataset has been generated using ESRI ArcMap software creating a GIS layer comprised of polygons that indicate the susceptibility to groundwater flooding. The susceptibility is split into three classes (A, B and C) of susceptibility to groundwater flooding. These are defined as follows:

Onshore areas outside of these classes are not considered to be prone to groundwater flooding based on the data used. The dataset does not show the likelihood of groundwater flooding occurring and does not take account of the chance of flooding from groundwater rebound. It covers a large area of land, and only isolated locations within the overall susceptible area are actually likely to suffer the consequences of groundwater flooding. The data should be used only in combination with other information, for example local data or historic data.

# NORTH DORSET DISTRICT COUNCIL LEVEL 1 SFRA

**APPENDIX A** 

**BGS** Susceptibility to Groundwater Flooding

Index Number: ND\_16

This document is the property of Jeremy Benn Associates Ltd. It shall not be reproduced in whole or in part, nor disclosed to a third party, without the permission of

